



# WARNING

KEEP OUT OF REACH OF CHILDREN

## POD-LOCK®

*Anti pod shatter sticker*

**For the reduction of pod-shatter in oilseed rape and other brassica seed crops**

### Active ingredient

SYNTHETIC LATEX 450 g/L

Also contains ALCOHOLS, C10-12, ETHOXYLATED PROPOXYLATED 50–100 g/L

**Net contents: 10 L**

Bar code to be inserted



**PLEASE READ THE LABEL COMPLETELY BEFORE USE**

**Any person purchasing, handling or disposing of this product must be suitably qualified.**



**WARNING**

Causes serious eye irritation.

Harmful to aquatic life with long lasting effects.

**PRECAUTIONARY STATEMENTS**

**PREVENTION**

Wash hands thoroughly after handling.

Wear eye protection/face protection.

Do not apply directly into or onto water.

Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area.

**RESPONSE**

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

If eye irritation persists: Get medical advice.

**DISPOSAL**

Dispose of contents/container in accordance with local regulations.

**Container disposal:** Ensure the container is empty. Triple rinse empty container and add rinsate to the spray tank. Recycle empty container through Agrecovery® (0800 247 326, [www.agrecovery.co.nz](http://www.agrecovery.co.nz)). Otherwise crush and bury in a suitable landfill. DO NOT reuse this container for any other purpose.

**Product disposal:** Dispose of this product only by using according to the label, or at an approved landfill or other approved facility.

**FIRST AID**

If swallowed do NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin contact occurs, remove contaminated clothing and flush skin with running water for several minutes.

**STORAGE**

Store in original container tightly closed and in a locked, dry, cool, well-ventilated area, away from food, seeds and foodstuffs. Protect from frost.

Storage must be generally in accordance with NZS8409 management of Agrichemicals. See Haznote for further information.

**SHELF LIFE**

When stored appropriately, this product should show no significant degradation for 2 years from the date of manufacture. When using this product beyond this shelf life, contact the registrant for further information.

**SAFETY DATA SHEET**

If additional hazard or safety information is required, refer to the Safety Data Sheet. For a copy visit our website at [www.upl-ltd.com/nz](http://www.upl-ltd.com/nz).

**EXCLUSION OF LIABILITY – PLEASE READ**

To the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, UPL New Zealand Limited as distributor, nor any other distributor have any liability or responsibility

whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

**Registered to:**  
**UPL New Zealand Ltd**  
PO Box 51584  
Pakuranga 2140  
Auckland  
Phone: 0800 100 325  
[www.upl-ltd.com/nz](http://www.upl-ltd.com/nz)

Registered pursuant to the ACVM Act 1997, No. P008857  
See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.  
Approved pursuant to the HSNO Act 1996. Approval Code HSR002503  
See [www.epa.govt.nz](http://www.epa.govt.nz) for approval controls.

© Copyright UPL New Zealand Limited 2020, all rights reserved.  
All directions for use, recommendations, use rates, compatibilities and other technical information is proprietary to UPL New Zealand Limited.

® Registered trademark

Batch number:	
Date of Manufacture:	

**In a transport emergency dial 111, Police or Fire Brigade.**  
**For specialist advice in an emergency only, call 0800 CHEMCALL (0800 243 622)**

**UN-FREE**

## GENERAL INFORMATION

**POD-LOCK** is a unique polymer blend which has been specifically developed to dramatically reduce the quantity of seed shed from seed pods splitting before and during harvest. For use in seed brassica crops including oilseed rape.

It does not slow down drying and has no adverse effects on maturity or quality of seed produced.

It should be applied between pods green and bendy stage (BBCH 80) until fully ripe – nearly all pods ripe, seeds black and hard (BBCH 89).

## DIRECTIONS FOR USE

CROP	USE RATE	COMMENTS
Brassica seed crops including oilseed rape	1 L/ha	Apply in 150-300 L/ha as a medium quality spray.  Use the 300 L/ha water volume on dense crops or for maximum effect.

For specific recommendations contact UPL New Zealand Limited or its distributors.

**It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.**

## MIXING

Three-quarter fill the spray tank with water and commence agitation. Add the required quantity of **POD-LOCK**. Continue agitation while topping up the tank and during spraying. May be tank mixed with glyphosate or diquat.

## EQUIPMENT

Immediately after application rinse the sprayer thoroughly with water and add **ACTIWETT**® or **RANGER**® non-ionic wetter or **ALL CLEAR**® **2X** tank decontaminant to clean **POD-LOCK** from the tank, pump, spray lines and nozzles.

® Registered trademarks